

RESEARCH ARTICLE

Open Access

Elevated arterial blood pressure and body mass index among Nigerian preschool children population

Odutola I Odetunde^{1*}, Emeka E Neboh⁴, Josephat M Chinawa², Henrietta U Okafor^{1,2,3,4}, Oluwatoyin A Odetunde⁵, Osita U Ezenwosu³ and Uchenna Ekwochi⁶

Abstract

Background: Arterial blood pressure tends to rise with growth and development. Elevated blood pressure (EBP) in children usually occurs during the first two decades of life, and the children with hypertension tend to grow into adulthood with the high level of blood pressure. The prevalence of hypertension in children is increasing, the causes likely to be of different combination of factors. In this study we ascertained the prevalence of EBP in pre-school children in Enugu metropolis, South-East Nigeria and also determined its association with some factors like the Body Mass Index (BMI), urinalysis finding, family history, gender, age and socioeconomic class.

Method: A Stratified method of sampling was used to select subjects from registered nursery schools (Pre- elementary school) within Enugu metropolis. Physical examination of the recruited pupils was done with emphasis on arterial blood pressure, anthropometric measurements and urinalysis.

Result: Six hundred and thirty children (630) were studied out of which 345 (54.8%) were males and 285 (45.2%) were females. Sustained EBP (mainly systolic) were recorded in 12 pupils (1.9%) giving a prevalence of 1.9% of the pre-school population. The twelve (1.9%) pupils were all 5 years of age (p value = 0.001) and 11 (1.72%) of them were of under-weight BMI. The prevalence of obesity is 0.5% and that of under-weight is 92% of the studied population. There is no association between EBP and obesity (p value = 0.679). All the pupils with EBP had protein-free urine and no hematuria.

Conclusions: EBP and under-weight malnutrition is common in children in 5 years age group. EBP in preschool children is not influenced by their body mass index, urinalysis finding, gender, family history of hypertension or socioeconomic class.

Keywords: Hypertension, Pre-hypertension, Body mass index (BMI), Preschool children, Nigeria

Background

The prevalence of hypertension in children is on the increase [1], as blood pressure increases with growth and development and results in hypertension during the first two decades of life [2]. Children with elevated blood pressure tend to maintain that level of blood pressure into adulthood [3]. Early detection with effective control is essential so as to minimize cardiovascular complications in adulthood. The risk factors for hypertension

includes obesity, family history of hypertension, change in dietary habits, decrease in physical activities and increasing stress as affected by the socioeconomic status [4]. The prevalence of hypertension in children varies between studies and depends on the studied age groups and the definition of hypertension in the studied population. Hypertension is defined as systolic or diastolic blood pressure greater than or equal to the 95th percentile for age and gender measured on at least three separate occasions and pre- hypertension as systolic or diastolic blood pressure between 90th and 95th percentile [5].

In children, systemic hypertension is uncommon with only about 1 percent of the childhood population having

* Correspondence: odetoyintola@yahoo.co.uk

¹Pediatric Nephrology Unit, Department of Pediatrics, University of Nigeria Teaching Hospital, UNTH, PMB 01129 Ituku-Ozalla Enugu State, Nigeria
Full list of author information is available at the end of the article

blood pressure significantly above the normal range [6]. Several studies on prevalence and risk factors for hypertension like high Body mass index (BMI) have focused more on older children and adolescents with less attention on the pre-school age children [7-11]. It is assumed that abnormal BMI in pre-school age children may not be related to medical problem and that health problem may not emerge until a child is overweight for several years (probably till adolescent age) before the evolution of obesity and its related health problem [12].

In this study we evaluated the usefulness of simply screening for high blood pressure and BMI calculation in early detection of blood pressure disorders in healthy pre-school children in Enugu metropolis in southeast Nigeria, with the objectives of determining the prevalence of pre hypertension and hypertension in these children and to ascertain its relationship with some known risk factors in asymptomatic subjects.

Methods

Setting

This study was carried out in Enugu metropolis, the capital city of Enugu State south-east of Nigeria. The population of the preschool age group in Enugu town was 37,556 (7 percent of total population) [13]. Written informed consent was obtained from the parents or the caregivers of the subjects before being enrolled into the study. The Ethical committee of the University of Nigeria Teaching Hospital (UNTH) gave approval for the study and permission was also obtained from State Ministry of Education before the commencement of the study.

Recruitment of subjects

A Stratified method of sampling was employed to get a sample that represents the population. The three local government areas (Enugu north, Enugu east and Enugu south) were used as the strata. In each stratum (local Government Area) four nursery schools were selected randomly from a constructed sampling frame of 75 Nursery schools.

In each of the nursery schools, a sample size of 50–60 pupils was calculated and were selected randomly based on population of the school and were given the proforma with universal sample bottle for urine collection. Thus, the numbers of proforma given per-school depend on the population of the school. Selection of schools was influenced by the cooperation and readiness of the management to participate in the research. Six hundred and thirty ($n = 630$) subjects gave their informed consent to be part of the study and also met the inclusion criteria of relatively healthy children for the study. Exclusion criteria include subjects with age less than 2 and greater than 5 years old (at the last birthday), presence of symptoms of renal disease like dysuria, increased urinary frequency or urgency,

macroscopic hematuria and facial or pedal edema, Presence of fever a week prior to the study, Lack of parental or guardian consent.

We carefully made sure that confounding and modifying factors of proteinuria and hematuria such as fever, intense activity or exercise, dehydration, emotional stress were ruled out in our study. This was done with thorough history taking and physical examination.

Methods

Physical examinations which included blood pressure, height, weight, pulse rate, and temperature measurements were carried-out on the subjects by trained physicians in the research team coordinated by the principal author in a standardized way. Emphasis of the physical examinations was on Body Mass Index (BMI) and Blood Pressure (BP) measurement. The BMI was calculated by the formula: $BMI = \text{Weight (Kg)}/\text{height (M}^2\text{)}$. We used the World Health Organization and Centers for Disease Control to classify BMI into underweight as BMI less than the 5th percentile, healthy weight as BMI of 5th up to the 85th percentile, overweight as BMI of 85th to less than the 95th percentile and obese as BMI equal to or greater than the 95th percentile for age and gender [14,15]. The blood pressure was measured by one observer. Seven or nine centimeter wrap-around cuffs were used depending on which width most closely approximated two thirds of the length of the upper arm. A mercury sphygmomanometer and an open-bell side of pediatrics-size Rappoport stethoscope were employed. Readings were recorded from right arm of the pupil in sitting position at the end of the routine examination and if the pupil was crying, the blood pressure was not taken until he or she became calm. The determined systolic and diastolic pressures were ascertained from three readings using the mean value of the three. An interval of 60 minutes was allowed and the cuff was completely deflated between readings. The systolic reading was taken as the first korotkoff-sound and the diastolic pressure was taken as the point where complete disappearance of the sound occurred (onset of 5th korotkoff-sound). We used the National High Blood Pressure In Children and Adolescent (2004 Working Group) guideline for diagnosis and Classification to classify the subjects into Normal BP with reading less than 90th percentile, Pre-hypertension with BP more than 90th percentile but less than 95th percentile while Hypertension was considered in subject with average reading greater than or equal to 95th percentile for age, gender, weight and height measured in at least three separate occasion [5].

We ensured stability of all our measurements through a constant reading of all our measurements that when other researchers measured same parameters, the error margin was insignificant. This is called “inter-rater reliability” or “inter-rater agreement [16]”.

Early morning urine specimens were collected from the subjects with sustained elevated BP and the completed proforma. The urine specimen was divided into 2 aliquots of 2 mls and 10 mls. The urinalysis was done on the uncentrifuged first aliquot (2 mls) of urine specimen using Combi 9 test strips by one of the authors. The second aliquot of the urine specimen and urine specimen of subjects with abnormal result (proteinuria and/or haematuria) were then transferred to the 10 ml test tubes, and were serially labeled and sent for microscopy. The specimens were centrifuged at 2000 rpm for 5 minutes. The microscopy was done on the sediments to determine presence of cast, red blood cells, white blood cells and crystals. From the proforma filled by the parents or care givers the family history of hypertension and renal disease was retrieved.

The families were assigned socioeconomic classes using the recommended method (modified) by Oyedeji [17]. The parents' occupation and highest education attained were scored from 1 (highest) to 5 (lowest). The mean score for both parents gives social class falling within the 1–5 range. Those with the mean score of < 2 were further reclassified into upper class while those with the mean score of > 2 were reclassified into lower social class. According to the protocol, for the occupation score, those in upper social class included parents, such as senior public officers, large-scale traders, large-scale farmers and professionals. Lower class included artisans, primary school teachers, peasant farmers, labourers and the unemployed. For the education score, those with PhD, master degree, bachelor degree and higher national diploma (HND) were categorized as upper class. Those with ordinary national diploma (OND), national certificate of education (NCE), technical education, grade II teachers' certificate, junior and senior secondary school certificate, primary school certificate and those with no formal education were classified as lower social class [17].

Statistical analysis

Data analysis was done using statistical package for social sciences (S.P.S.S for windows 17.0 output). Frequencies were compared using Chi squared test. Chi square was also used to test for association between differences in proportions of the independent variables (BMI, Age, Gender, Urine analysis findings and Family background) on the dependent variables (Blood Pressure reading). All the continuous variables were re-grouped into categorical forms (e.g. BMI was regrouped as Under Weight, Healthy Weight, Over-Weight and Obese) also, age has its categories as age two, age three, age four and five. Sex was grouped as Male and female, other categorical variables include family history of blood pressure and family history of renal diseases grouped as **YES** for those with history of renal diseases and high blood pressure in their family, and **NO** for those who have no history of renal diseases and

high blood pressure in their family. P value of < 0.05 was considered significant. Data presentation was in tables.

Results

A total of 630 subjects were studied comprising of 345 males (54.8%) and 285 females (45.2%), giving a male: female ratio of 1: 0.8. Of the 345 male subjects, 66 (19.1 percent) were aged 2 years; 85 (24.7%) 3 years; 95 (27%) 4 years and 99 (28.7%) 5 years. On the other hand, of 285 female subjects, 55 (17.5%) were aged 2 years; 84 (29.5%) aged 3 years; 68 (23.9%) aged 4 years and 83 (29.1%) aged 5 years. Sustained elevated BP were recorded in 12 (1.9%) of the studied population nine (1.4%) of which had pre -hypertension and 3 (0.5%) hypertension (Table 1). The twelve (1.9%) subjects with elevated BP were aged 5 years (P value = 0.001) (Table 1). Of the twelve subjects

Table 1 Showing percentage distribution of children by the level of blood pressure in relation to their age, gender and BMI

Percentage distribution of the children by the level of their blood pressure				
Blood pressure (BP)	Frequency (N)	Percentage (%)		
Hypertension	3	0.5		
Pre- Hypertension	9	1.4		
Normal BP	618	98.1		
Total	630	100		

Distribution of elevated blood pressure by age of the children				
Age (Yrs)	Pre HBP	HBP	Normal	Total
2 - < 3	0 (0)	0 (0)	116 (18.4)	116 (18.4)
3- < 4	0 (0)	0 (0)	169 (26.8)	169 (26.8)
4 - < 5	0 (0)	0 (0)	163 (25.9)	163 (25.9)
5	9 (1.4)	3 (0.5)	170 (27.0)	182 (28.9%)
Total	9 (1.4)	3 (0.5)	618 (98.1)	630 (100)

$\chi^2 = 28.76$ df = 5 P- value = 0.001

Distribution of elevated blood pressure by gender of the children				
Gender	Pre HBP	HBP	Normal	Total
Male	6 (0.95)	2 (0.32)	337 (53.5%)	345 (54.8)
Female	3 (0.45)	1 (0.18)	281 (44.6)	285 (45.2)
Total	9 (1.4)	3 (0.5)	618 (98.1)	630 (100)

$\chi^2 = 0.70$ df = 2 P-value = 0.71

Distribution of elevated blood pressure by BMI of the children				
BMI Classification	Pre HBP	HBP	Normal	Total
Underweight	9 (1.4)	2 (0.32)	570 (90.5)	581 (92.2)
Healthy weight	0 (0)	1 (0.18)	43 (6.8)	44 (7)
Overweight	0 (0)	0 (0)	2 (0.3)	2 (0.3)
Obese	0 (0)	0 (0)	3 (0.5)	3 (0.5)
Total	9 (1.4)	3 (0.5)	618 (98.1)	630 (100)

$\chi^2 = 3.98$ df = 5 P- value = 0.68

with elevated blood pressure 8 (1.3%) were male while other 4 (0.6%) were females (Table 1). Three (0.5%) of the subjects were Obese, 2 (0.3%) were over-weight while 581 (92.2%) were underweight. Elevated BP has no association with obesity or over-weight (Table 1). The variation of result within the social classes was not significant (P value = 0.56) (Table 2).

There is positive family of hypertension in 71 (11.3%) of the subjects out of which only 2 (0.3%) had elevated BP (Table 3). None of the subjects with elevated BP had positive family history of renal disease.

All the subjects with elevated BP had protein-free urine and no hematuria. Hypertension was observed in all the social classes of the studied population.

Discussion

There are several technical challenges in comparing various prevalence studies on elevated arterial BP in children because of the dissimilarity in the studied population and criteria used in defining the condition. In this study, we employed the definition of elevated BP based on the 2004 National High Blood Pressure Education Program Working Group in Children and Adolescent to evaluate and classify our subjects [6]. Our studied population was healthy preschool age children in which the prevalence of elevated BP (systolic) of 1.9% was obtained, 1.4% and 0.5% for pre-hypertension and hypertension respectively. There are paucity of studies on elevated BP and risk factors in the preschool age children [18,19]. The prevalence of elevated BP in this study is lower when compare with 5.2% and 13% obtained by Vitolo et al. and William et al. respectively in their studies of a similar age population though in different settings [18,19]. The prevalence of 1.9 percent in this study is slightly higher when compared with that of 1% by National High Blood Pressure Education Program Working Group in Children in the United State of America [6]. In this study, subjects with hypertension were limited to children in 5 years age group with no gender preponderance. This is similar to the observation of Still and Cottom [20] who noted higher incidence of hypertension in the age group 5 to 10 years with no gender preponderance. The study was carried out in a

Table 2 Distribution of subjects with elevated blood pressure by socio-economic class

Class	Blood pressure		Total n (%)
	High n (%)	Normal n (%)	
Upper	2 (2.1%)	93 (97.9)	95 (100)
Middle	7 (2.0%)	436 (98.0)	443 (100)
Lower	3 (3.2%)	89 (96.8)	92 (100)
Total	12 (1.9)	618 (98.1)	630 (100)
$\chi^2 = 1.18$ df = 2 P-value = 0.56.			

Table 3 Family history of high blood pressure and renal disease of the children by blood pressure

Family history of Hypertension	Blood pressure			TOTAL n (%)
	Pre HBP n (%)	HBP n(%)	NORMAL n (%)	
Yes	1 (0.18)	1 (0.18)	69 (11)	71 (11.3)
No	8 (1.22)	2 (0.32)	549 (87.1)	559 (88.7)
TOTAL	9 (1.4)	3 (0.5)	618 (98.1)	630 (100)
$\chi^2 = 1.5$ df = 2 P-value = 0.5				
Family history of Renal disease	Blood pressure			TOTAL n (%)
	Pre HBP n (%)	HBP n (%)	NORMAL n (%)	
Yes				
No	8 (1.22)	2 (0.32)	549 (87.1)	559 (88.7)
TOTAL	9 (1.4)	3 (0.5)	618 (98.1)	630 (100)
$\chi^2 = 1.5$ df = 2 P-value = 0.5				

metropolitan city where due to the challenges of low acceptability of exclusive breastfeeding initiative, high acceptability of breast milk substitutes and modernization of diet in the general population, majority (92.2%) of the subjects were underweight while only 7% had healthy weight.

The prevalence of obesity and overweight were 0.5% and 0.3% respectively and there were no association between these and elevated BP. Several studies in the general population in Nigeria have documented prevalence rate of obesity in children and adolescents to range from 0.3 – 18% [21-23] and Rosnner et al. in the study of 47,000 children reported that the risk of elevated blood pressure is higher in children in upper deciles of BMI compared with odd ratio of systemic hypertension ranging from 2.5 -5.7 mmHg [24]. Our finding of prevalence for obesity and overweight are in contrast with the report by Williams et al. [18] in a similar population though in an industrialized country where they documented a prevalence of 15% and 17% for obesity and overweight respectively while 13% had elevated BP. Virtually all the subjects with EBP in our study were of underweight BMI. This is in consonant with the findings of Sawaya AL et al. [25] in their study done in another developing country. They reported that the prevalence of hypertension in under-nourished preschool children and those that recovered from under nutrition to be higher than that of the controls with normal nutrition and concluded that this may possibly impact on the morbidity and mortality related to hypertension during adulthood. Nearly all our subjects (92.2%) had underweight BMI. This may be a reflection of nutritional deficiency during prenatal life that continues in the post-natal period which is very common in the developing countries [25].

In this study, no association was observed between hypertension, proteinuria or hematuria which are surrogate

markers for renal disease. The distribution of proteinuria and hypertension in this study had no significant preponderance for any particular socioeconomic group, although some of the renal disorders in Africa are related to infection and infestation which are commoner in the lower social class. There were also no significant association between EBP and positive family history of hypertension or renal disease in the subjects. Several studies had reported positive family history of hypertension, renal disease, change in dietary habits, decrease physical activities and increasing stress as affected by the socioeconomic status as risk factors for development of EBP in children [4,18,19].

Conclusions

There is a significant association between EBP and pupils in 5 years age group while obesity, urinalysis finding, gender, family history and socioeconomic class do not affect BP.

Pre-admission health examination should be introduced at primary school level and should include BP and BMI measurement in order to identify those with EBP. All cases of elevated blood pressure with or without abnormal BMI detected during this screening should be referred to Pediatric Nephrologists for proper evaluation and follow-up.

Abbreviations

BMI: Body mass index; EBP: Elevated blood pressure.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

OIO, EEN, JMC and OUE envision the study, participated in its design, data collection and coordination of the study and drafted the manuscript. HUO, JMC, OAO and UE contributed substantially to data collection, analysis and interpretation and revision of the manuscript. OIO, EEN, OAO, HUO, JMC, UE and OUE did the final revision of the manuscript. All authors read and approved the final manuscript for publication.

Acknowledgement

We wish to acknowledge the International Pediatric Nephrology Association and International Society of Nephrology for giving the principal author the opportunity to undergo fellowship training programme through their sponsorship. The authors also expressed their profound gratitude to Mr Peter Okunade a research statistician, for his advice on the analysis of data. Finally, our special thanks go to Mrs. Deola Ogunwole, a former staff nurse with U.N. T.H Nursing services, Dr. Bosah Umerah, Uche Azuike for being part of the research team in data collection and all the proprietors of the Nursery schools used for the study for the assistance rendered towards the successful completion of this study.

Author details

¹Pediatric Nephrology Unit, Department of Pediatrics, University of Nigeria Teaching Hospital, UNTH, PMB 01129 Ituku-Ozalla Enugu State, Nigeria. ²Cardiology Unit, University of Nigeria Teaching Hospital, UNTH, PMB 01129, Ituku-Ozalla, Enugu State, Nigeria. ³Oncology and Hematology Unit, University of Nigeria Teaching Hospital, UNTH, PMB 01129, Ituku-Ozalla, Enugu State, Nigeria. ⁴Department of Medical Biochemistry, College of Medicine, Enugu State University of Science and Technology (ESUT), Enugu, Nigeria. ⁵Pediatric Surgery Unit, Department of Surgery, Enugu State

University Teaching Hospital, Enugu, Nigeria. ⁶Department of Pediatrics, Enugu State University Teaching Hospital, Enugu, Nigeria.

Received: 20 June 2013 Accepted: 26 February 2014

Published: 4 March 2014

References

1. Din- Dzietham R, Liu Y, Bielo MV, Shamsa F: High blood pressure trends in children and adolescents in national surveys. 1963 to 2002. *Circulation* 2007, **116**:1488-1496.
2. Chadha SL, Tandon R, Shekhawat S, Gopinath N: An epidemiology study of blood pressure in school children (5-14 years) in Delhi. *Indian Heart J* 1999, **51**:178-182.
3. Anjana P, Kaur N, Kumari K, Sidhu S: Variation in blood pressure among school children of Amritsar (Punjab). *Anthropologist* 2005, **7**:201-204.
4. Mohan B, Kumar N, Aslam N, Rangbulla A, Kumbkarni S, Sood NK, Wander GS: Prevalence of sustained hypertension and obesity in urban and rural school going children in Ludhiana. *Indian Heart J* 2004, **56**:310-314.
5. Chobanian AV, Bakris GL, Black HR, Cushman WC, Green LA, Izzo JL Jr, Jones DW, Materson BJ, Oparil S, Wright JT Jr, Roccella EJ: The seventh report of the joint national committee on prevention, detection, evaluation, and treatment of high blood pressure. *Hypertension* 2003, **42**:1206-1252.
6. National high blood pressure education program working group. Update on the 1987 task force report on high blood pressure in children and adolescent: A working group report from the national high blood pressure education programme. *Paediatrics* 1996, **98**:649-658.
7. Mo-suwan L, Junjana C, Puetpaiboon A: Increasing obesity in school children in transitional society and the effect of the weight control programme. *Southeast Asian J Trop Med Public Health* 1993, **24**:590-594.
8. Gupta AK, Ahmad AJ: Childhood obesity and hypertension. *Indian Pediatr* 1990, **27**:333-337.
9. Bugage MA, Yakubu AM, Ogala WW: Prevalence of adolescent hypertension in Zaria. *Nig J Paediatr* 2005, **32**:77-82.
10. Antia Obong OE, Anita Obong IE: Arterial blood pressure of Nigeria urban- rural school children. *Nig J Paediatr* 1991, **18**:3-11.
11. Ejike CECC, Ugwu EC, Ezeanyika LUS: Variation in the prevalence of (pre) hypertension in a Nigerian school- going adolescent population living in a semi-urban and an urban area. *BMC Pediatr* 2011, **10**:10-13.
12. Hernandez B, Uphold CR, Graham MV, Singer L: Prevalence and correlation of obesity in preschool children. *J Pediatr Nurs* 1998, **13**:68-78.
13. Nigeria 2006 Census figure: <http://www.nigeriamasterweb.com/Nigeria06CensusFigs.html>. last accessed on 2013 June 12.
14. BMI Percentile Calculator for Child and Teen, English Version. CDC Home: <http://apps.nccd.cdc.gov/dnpabmi/Calculator.aspx>. Last assessed on 2013 June 14.
15. Kuczmarki RJ, Ogden CL, Grummer-Strawn LM, Flegal KM, Guo SS, Wei R, Mei Z, Curtin LR, Rocche AF, Johnson CL: CDC growth charts: United State. *Adv Data* 2000, **314**:1-27.
16. Geeta A, Jamaiah H, Safiza MN, Khor GL, Kee CC, Ahmad AZ, Suzana S, Rahmah R, Faudzi A: Reliability, technical error of measurements and validity of instruments for nutritional status assessment of adults in Malaysia. *Singapore Med j* 2009, **50**:1013-1018.
17. Oyediji GA: Socio- economic and cultural background of hospitalized children in Ilesha. *Nig J Pediatr* 1985, **12**:111-117.
18. William CL, Strobino B, Bolllella M, Brotanek J: Body size and cardiovascular risk factors in preschool population. *Prev Cardiol* 2004, **7**:116-121.
19. Vitolo MR, da Costa Louzada ML, Rauber F, Campagnolo PD: Risk factors for high blood pressure in low income children aged 3-4 years. *Eur J Pediatr* 2013. doi:10.1007/s00431-013-2012-9.
20. Still JL, Cotton D: Severe hypertension in childhood. *Arch Dis Child* 1967, **42**:34-39.
21. Okesina AB, Oparinde DP, Akindoyin KA, Erasmus RT: Prevalence of some risk factors of coronary heart disease in rural Nigeria population. *East Afr Med J* 1999, **76**:212-216.
22. Ammen IO, Fawole OI: Prevalence of cardiovascular disease among the youths. Investing in the young people health and development research that improves policies and programmes. *Abuja Nigeria* 2008. Online www.jhsph.edu/gatesinstitute/pdf/policypractice. last accessed 2013, June 14.
23. Owa JA, Adejuyigbe O: Fat mass percentage body mass index and mid-upper arm circumference in a healthy population of Nigerian children. *J Trop Pediatr* 1997, **43**:13-19.

24. Rosnar B, Prineas R, Daniel SR: **Blood pressure difference between black and white in relation to body size among US children and adolescents.** *Am J Epidemiol* 2000, **5**:1007–1019.
25. Sawaya AL, Sesso R, Florencio TM, Fernandes MT, Martins PA: **Association between chronic under nutrition and hypertension.** *Matern Child Nutr* 2005, **1**:155–163.

doi:10.1186/1471-2431-14-64

Cite this article as: Odetunde et al.: Elevated arterial blood pressure and body mass index among Nigerian preschool children population. *BMC Pediatrics* 2014 **14**:64.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

