

RESEARCH

Open Access



# Factors associated with in-hospital recurrence of intestinal intussusception in children

Jing Zhang<sup>1</sup>, Qi Dong<sup>1\*</sup>, Xiaoxia Su<sup>1</sup> and Junshan Long<sup>1</sup>

## Abstract

**Background** A minority of children experience in-hospital recurrence of intestinal intussusception after treatment. This study investigated the factors associated with in-hospital recurrence of intussusception in pediatric patients in China.

**Methods** This retrospective study included patients aged 0–18 years-old with intestinal intussusception treated at Hainan Women and Children's Medical Center between January 2019 and December 2019. Demographic and clinical characteristics were extracted from the medical records. Factors associated with in-hospital recurrence of intussusception were identified by logistic regression analysis.

**Results** The analysis included 624 children (400 boys) with a median age of 1.8 years (range, 2 months and 6 days to 9 years). Seventy-three children (11.7%) had in-hospital recurrence of intussusception after successful reduction with air enema. Multivariate logistic regression analysis identified age > 1 year-old (odds ratio [OR]: 7.65; 95% confidence interval [95%CI]: 2.70–21.71;  $P < 0.001$ ), secondary intestinal intussusception (OR: 14.40; 95%CI: 4.31–48.14;  $P < 0.001$ ) and mesenteric lymph node enlargement (OR: 1.90; 95%CI: 1.13–3.18;  $P = 0.015$ ) as factors independently associated with in-hospital recurrence of intussusception.

**Conclusions** Age > 1 year-old, secondary intussusception and mesenteric lymph node enlargement were independently associated with increased odds of in-hospital recurrence of intussusception after successful reduction with air enema.

**Keywords** Intussusception, Intestinal, Infant, Child, Risk factors, Recurrence

## Background

Intussusception describes the intussusception of a segment of intestine into the lumen of a more distant segment of the bowel and is one of the most common acute abdominal disorders in infancy [1]. Intestinal intussusception occurs most frequently at the age of 6 months [1], and the incidence of intussusception in China during

2009–2013 was estimated to be 112.9 per 100,000 children aged < 2 years-old [2]. Intussusception in children is a medical emergency that necessitates timely diagnosis and treatment [3]. The main presenting symptoms of intestinal intussusception in children under 2 years of age include vomiting, abdominal pain and blood in the stools [4]. The diagnosis of intussusception is most often made using ultrasonography, but other useful investigations include X-ray, computed tomography and barium/air enema [1, 5, 6]. Management strategies include reduction by pneumatic (air) enema, hydrostatic (liquid) enema, laparoscopic surgery and open surgery [3, 7, 8].

\*Correspondence:

Qi Dong  
dqearth@163.com

<sup>1</sup>Department of General Surgery, Hainan Women and Children's Medical Center, Haikou 570100, China



© The Author(s) 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Previous research has indicated that a recurrence of intussusception occurs in 8–19% of children [9–13]. Several clinical analyses have attempted to identify factors associated with an increased risk of intussusception recurrence. Many investigations have concluded that older age ( $\geq 1$  year-old or  $\geq 2$  years-old, depending on the analysis) is associated with elevated odds of intussusception recurrence [13–16]. Pathological lead points (PLPs), such as Meckel's diverticulum, duplication, polyps or tumors, have also been reported as independent predictors of recurrent intussusception [13, 14, 16]. Other factors reported to increase the risk of intussusception recurrence include an absence of vomiting [13], rectal bleeding [16], fever [17], surgical treatment [18], overweight/obesity and white blood cell (WBC) count  $\geq 20 \times 10^9 / L$  [19]. However, there have also been disagreements between studies. For example, Guo et al. found that symptom duration  $\leq 12$  h and mass located in the right abdomen were independent predictors of recurrent intussusception [13], whereas the factors identified by Xie et al. included duration of symptoms  $\geq 48$  h and mass located in the left abdomen [16]. Furthermore, Zhang et al. demonstrated that mass diameter  $> 2.55$  cm and enlarged abdominal lymph nodes were related to recurrence of intussusception [20], whereas Vo et al. did not identify any radiologic features that were associated with early recurrence, although the risk of recurrence was higher in females and patients with fever [17]. A recent meta-analysis by Ye et al. identified only fever, absence of vomiting and PLPs as factors associated with a higher risk of recurrence of intestinal intussusception [21].

At least half of all recurrences of intussusception occur within 48 h after treatment [14, 17, 22, 23], with a median time to recurrence of 25 h after fluoroscopy-guided air enema [15]. Furthermore, Lee et al. reported that in-hospital recurrence of intussusception occurred in 6.1% of pediatric patients [24]. Knowledge of the risk factors for early recurrence of intestinal intussusception would help pediatric surgeons to identify those patients requiring careful observation during the early post-operative period. However, only a relatively small number of studies have evaluated the clinical factors associated with an increased risk of intussusception recurrence, but the findings have varied. Simanovsky et al. found that patients with early recurrence were older than those who did not have early recurrence (23 months-old vs. 12 months-old) [25], while Shen et al. concluded that age  $\geq 2$  years-old and absence of fever were associated with short-term recurrence [26]. Another analysis highlighted higher body weight, PLPs and onset during the second quarter of the year as factors predictive of early recurrence [27]. Additionally, a recent study by Zhu et al. showed that a blood level of monocyte chemotactic protein-1 exceeding

23.24 ng/mL predicted short-term recurrence of intussusception with a sensitivity of 82.1% and a specificity of 75.7% [28]. However, Cho et al. did not find any differences in age, reduction success rate, treatment method (surgery vs. non-surgery) or PLPs between patients with recurrence within 48 h and those without recurrence [29]. Thus, the findings of these studies are not entirely consistent.

The aim of this study was to investigate the factors associated with in-hospital recurrence of intestinal intussusception in pediatric patients in China. To achieve this aim, we performed a retrospective analysis of data for patients aged  $\leq 18$  years-old who were treated for intussusception at our institution over a 12-month period.

## Methods

### Study design and patients

This retrospective study included patients with intestinal intussusception treated at Hainan Women and Children's Medical Center (Haikou, Hainan, China) between January 2019 and December 2019. The inclusion criteria were as follows: (1) aged 0–18 years-old; (2) the diagnosis of intussusception was confirmed by air enema imaging; and (3) intussusception was successfully reduced by air enema. Intussusception was defined as intestinal obstruction caused by the insertion of a segment of intestine and its corresponding mesentery into the adjacent intestinal cavity. There were no formal exclusion criteria. The study protocol was approved by the Ethics Committee of Hainan Women and Children's Medical Center (HNWCMC Tribunal No. [26], 2023). The requirement for informed consent was waived because the analysis was retrospective and anonymized.

### Data collection and definitions

The following demographic, clinical and ultrasonographic characteristics were obtained from the medical record system: age, sex, symptoms (including fever, vomiting and bloody stools), duration of symptoms prior to admission, history of previous intussusception, type of intestinal intussusception (primary [idiopathic] or secondary to an identifiable organic lesion), site of the lesion (ileocecal junction, ascending colon, hepatic flexure, transverse colon, splenic flexure or other), lesion size (inner diameter and outer diameter), mesenteric lymph node (enlargement or normal), C-reactive protein (CRP) level, WBC count, neutrophil percentage and lymphocyte percentage. The normal mesenteric lymph node size of the general pediatric population was 0.2–0.5 cm. There were more than 1 lymph node in the mesentery in the same area, with a short axis diameter  $> 0.5$  cm, long axis diameter  $> 1.0$  cm or aspect ratio  $> 2$ , or lymph nodes arranged in clusters, Color Doppler flow imaging showing increased blood flow in lymph nodes, were

considered mesenteric lymph node enlargement. Recurrence of intussusception specifically referred to the recurrence of intussusception during hospitalization.

### Statistical analysis

SPSS 26.0 software was used for the analysis. Normally distributed continuous data are described as the mean  $\pm$  standard deviation (SD) and were compared between groups using Student's *t*-test for independent samples. Non-normally distributed continuous data are presented as median (interquartile range [IQR]) and were analyzed using the Mann-Whitney *U* test. Categorical data are described as *n* (%) and were compared between groups using the chi-squared test or Fisher's exact test. Logistic regression analysis was used to identify factors associated with in-hospital recurrence of intussusception, and variables with statistical significance in the univariate analysis were included in the multivariate analysis. Odds ratios (ORs) and 95% confidence intervals

(95% CIs) were calculated. A two-sided *P*-value  $< 0.05$  was considered statistically significant.

## Results

### Baseline characteristics of the study participants

The analysis included 624 children (400 boys, 64.1%) with a median age of 1.8 years (range, 2 months and 6 days to 9 years). The age at presentation was  $\leq 1$  year-old in 178 children (28.5%),  $> 1$  year-old in 446 children (71.5%), among them,  $\geq 2$  years-old in 255 children (40.9%). Seventy-four children (11.9%) had a history of previous intestinal intussusception. Seventy-three children (11.7%) had in-hospital recurrence of intestinal intussusception after successful reduction with air enema. The baseline characteristics of the study participants are shown in Table 1. There were no significant differences between the in-hospital recurrence group and non-recurrence group in sex, history of previous intussusception, symptom profile, duration of symptoms, site of the intestinal intussusception, mass size, CRP level, WBC count, neutrophil

**Table 1** Baseline clinical characteristics of the study participants

Characteristic	All ( <i>n</i> = 624)	No recurrence ( <i>n</i> = 551)	Recurrence ( <i>n</i> = 73)	<i>P</i>
Age (years)	1.8 (1.00–2.8)	1.7 (0.9–2.7)	2.1 (1.7–3.3)	$< 0.001$
$\leq 1$ year-old	178 (28.5%)	174 (31.6%)	4 (5.5%)	0.001
$> 1$ year-old	446 (71.5%)	377 (68.4%)	69 (94.5%)	
Male	400 (64.1%)	346 (62.8%)	54 (74.0%)	0.061
Symptoms				
Fever	152 (24.4%)	134 (24.3%)	18 (24.7%)	0.950
Vomiting	333 (53.4%)	295 (53.5%)	38 (52.1%)	0.810
Bloody stools	120 (19.2%)	112 (20.3%)	8 (10.9%)	0.056
History of previous intussusception	74 (11.9%)	61 (11.1%)	13 (17.8%)	0.094
Type of intussusception				$< 0.001$
Primary	610 (97.8%)	546 (99.1%)	64 (87.7%)	
Secondary	14 (2.2%)	5 (0.9%)	9 (12.3%)	
Site of intussusception				0.806
Hepatic flexure	237 (38.0%)	206 (37.4%)	31 (42.5%)	
Transverse colon	184 (29.5%)	162 (29.4%)	22 (30.1%)	
Ileocecal junction	92 (14.7%)	84 (15.2%)	8 (11.0%)	
Ascending colon	63 (10.1%)	55 (10.0%)	8 (11.0%)	
Splenic flexure	31 (5.0%)	30 (5.4%)	1 (1.4%)	
Other	17 (2.7%)	14 (2.5%)	3 (4.1%)	
Duration of symptoms				0.901
$\leq 2$ days	575 (92.1%)	508 (92.2%)	67 (91.8%)	
$> 2$ days	49 (7.9%)	43 (7.8%)	6 (8.2%)	
Mesenteric lymph node enlargement	251 (40.2%)	210 (38.1%)	41 (56.2%)	0.003
Size of mass (cm)				
Inner diameter	2.8 (2.6–3.1)	2.8 (2.6–3.1)	2.9 (2.5–3.2)	0.220
Outer diameter	3.5 (3.1–3.8)	3.5 $\pm$ 0.7	3.6 $\pm$ 0.8	0.290
CRP (mg/L)	5.1 (1.8–11.4)	5.2 (1.8–11.0)	4.3 (1.8–13.0)	0.907
WBC count ( $\times 10^9/L$ )	11.0 (8.6–13.6)	10.9 (8.6–13.4)	11.1 (8.6–14.4)	0.655
Neutrophils (%)	62.4 (48.7–73.6)	61.3 $\pm$ 15.9	62.3 (46.6–70.9)	0.386
Lymphocytes (%)	30.1 (19.9–41.5)	30.8 $\pm$ 13.9	31.6 $\pm$ 13.2	0.544

Data are presented as median (IQR), mean  $\pm$  SD or *n* (%)

percentage or lymphocyte percentage (Table 1). However, patients with in-hospital recurrence of intussusception were older (median [IQR]: 2.08 [1.67–3.25] vs. 1.67 [0.92–2.67] years-old,  $P < 0.001$ ) and had significantly higher incidences of secondary intussusception (12.3% vs. 0.9%,  $P < 0.001$ ) and mesenteric lymph node enlargement (56.2% vs. 38.1%,  $P = 0.003$ ) than those who did not have recurrence.

#### Logistic regression analysis of factors associated with in-hospital recurrence of intussusception

The multivariate analysis identified age > 1 year-old (OR = 7.65, 95%CI = 2.70–21.71,  $P < 0.001$ ), secondary intestinal intussusception (OR = 14.40, 95%CI = 4.31–48.14,  $P < 0.001$ ) and mesenteric lymph node enlargement (OR = 1.90, 95%CI = 1.13–3.18,  $P = 0.015$ ) as factors independently associated with in-hospital recurrence of intussusception (Table 2).

#### Discussion

The main findings of this retrospective analysis were that age > 1 year-old, secondary intestinal intussusception and mesenteric lymph node enlargement were independently associated with increased odds of in-hospital recurrence of intussusception. We suggest that children with intestinal intussusception who have any of these clinical characteristics should be carefully monitored throughout their hospital stay to ensure that any recurrence of intestinal intussusception after treatment is diagnosed and managed in a timely manner.

In-hospital recurrence of intestinal intussusception occurred in 11.7% of the children included in this study. The incidence in our cohort agrees well with the overall rate of 8–19% described previously [9–13], given that over half of all recurrences occur within 48 h after treatment [14, 17, 22, 23]. The incidence of 11.7% is also broadly consistent with values reported by studies of early recurrence, including those by Lee et al. (in-hospital recurrence of 6.1%) [24], Zhu et al. (9.0% within 72 h of reduction) [28], Shen et al. (12.8% within 72 h of reduction) [27] and Adhikari et al. (16.7% within 1 month of treatment) [30].

**Table 2** Multivariate logistic regression analysis of factors associated with in-hospital recurrence of intestinal intussusception

Variable	Odds ratio (95% confidence interval)	P
Age (> 1 years-old vs. ≤ 1 years-old)	7.65 (2.70–21.71)	< 0.001
Type of intussusception (secondary vs. primary)	14.40 (4.31–48.14)	< 0.001
Mesenteric lymph node enlargement (yes vs. no)	1.90 (1.13–3.18)	0.015

The present study found that age > 1 year-old was an independent factor strongly associated with in-hospital recurrence of intussusception (OR = 7.65). This result agrees well with several published studies. For example, Guo et al. and Chen et al. identified age > 1 year-old as an independent predictor of the recurrence of intestinal intussusception [13, 14]. Similarly, Kim et al. and Xie et al. reported that age ≥ 2 years-old was independently associated with recurrent intussusception [15, 16]. Although the above four studies did not focus on in-hospital or early recurrence, other investigations specifically evaluating the early recurrence of intussusception after treatment have also identified older age as a risk factor. Simanovsky et al. concluded that children with recurrence of intussusception within 48 h of treatment were older than those who did not have early recurrence (23 months-old vs. 12 months-old) [25], while Shen et al. concluded that age ≥ 2 years-old was related to short-term recurrence [26]. However, Cho et al. did not find any differences in age between patients with recurrence of intussusception within 48 h and those without recurrence [29], and a similar finding was reported by Adhikari et al., who evaluated short-term recurrence within 1 month of treatment [30]. The reasons for the disparity between studies remain unclear, although it is possible that differences in analytical approach (e.g., grouping on the basis of < 1 year-old vs. < 2 years-old) may have contributed. In addition, given that the incidence of recurrence is quite low, it cannot be excluded that some studies were underpowered to detect a real association between age and recurrence of intestinal intussusception.

Secondary intestinal intussusception (i.e., intussusception due to the presence of identifiable organic disease) was strongly associated with disease recurrence in the present study (OR = 14.40). This finding concurs with several prior investigations demonstrating that PLPs are independent predictors of recurrent intussusception [13, 14, 16, 21, 27]. PLPs are associated with 0.3–20% of cases of intestinal intussusception in children [31, 32], with the main causative PLPs being Meckel's diverticulum, ileal polyp, duplication of the ileum, Peutz–Jeghers syndrome, malignant tumors and appendiceal stump [32]. It is becoming increasingly accepted that PLPs elevate the risk for intussusception recurrence [8], and the present study provides further evidence to support this.

Our analysis also identified mesenteric lymph node enlargement as a predictor of in-hospital recurrence of intestinal intussusception (OR = 1.90). This result agrees with the study of Zhang et al., who found that enlarged abdominal lymph nodes were related to recurrence of intussusception [20]. Adhikari et al. also observed that the incidence of lymph node enlargement was numerically higher in the recurrence group than in the non-recurrence group (61.5% vs. 47.7%), although the

difference was not statistically significant [30]. Enlarged lymph nodes have been reported to not only decrease the hydrostatic reduction rate but also act as a lead point in intussusception [33]. Thus, it is possible that the presence of mesenteric lymph node enlargement might act as a PLP to increase the risk of the intussusception reforming after initially successful treatment.

Among the other factors evaluated in this study, an absence of bloody stools ( $P=0.056$ ) and male sex ( $P=0.061$ ) were borderline significantly more common in the recurrence group than in the non-recurrence group. With regard to the former, Lee et al. also reported that bloody stools were less common in the recurrence group than in the non-recurrence group [24], whereas Xie et al. conversely found that rectal bleeding was a risk factor for recurrence [16]. With regard to the latter, most studies have not detected an influence of sex on intussusception recurrence, although one analysis suggested a higher risk of recurrence in females [17]. Thus, it seems unlikely that sex and bloody stools are major factors influencing the risk of recurrence of intestinal intussusception.

This study has some limitations. First, this was a retrospective analysis, so the results may have been affected by information bias or selection bias. Second, this was a single-center study, so the generalizability of the findings cannot be confirmed. Third, since the number of patients with recurrence of intussusception was quite small (due to the low risk of recurrence), the study may have been underpowered to detect some real differences between the recurrence and non-recurrence groups. Fourth, other unknown factors not included in the analysis may have influenced the results. Large-scale, prospective, multi-center studies are needed to further evaluate the factors associated with in-hospital recurrence of intestinal intussusception.

## Conclusions

In conclusion, this single-center, retrospective analysis of children treated for intestinal intussusception found that age > 1 year-old, secondary intestinal intussusception and mesenteric lymph node enlargement were independently associated with increased odds of in-hospital recurrence. Children with intestinal intussusception who have any of these clinical characteristics may require close observation during hospitalization to ensure that any recurrence of intussusception is diagnosed and treated in a timely manner.

## Abbreviations

PLPs	Pathological lead points
WBC	white blood cell
CRP	C-reactive protein
SD	standard deviation
ORs	Odds ratios

## Acknowledgements

Not applicable.

## Authors' contributions

JZ and QD carried out the studies, participated in collecting data, and drafted the manuscript. JSL and XXS performed the statistical analysis and participated in its design. JZ and QD participated in acquisition, analysis, or interpretation of data and draft the manuscript. All authors read and approved the final manuscript.

## Funding

This research was supported by Hainan province Clinical Medical Center (QWYH202175).

## Data availability

All data generated or analyzed during this study are included in this published article.

## Declarations

### Ethics approval and consent to participate

The study protocol was approved by the Ethics Committee of Hainan Women and Children's Medical Center (HNWCMC Tribunal No. [26], 2023). The requirement for informed consent was waived by the ethics committee of Hainan Women and Children's Medical Center because of the retrospective nature of the study. All methods were conducted in accordance with relevant guidelines and regulations. All procedures were performed in accordance with the ethical standards laid down in the 1964 Declaration of Helsinki and its later amendments.

### Consent for publication

Not applicable.

### Competing interests

The authors declare no competing interests.

Received: 6 April 2023 / Accepted: 23 August 2023

Published online: 26 August 2023

## References

1. Jiang J, Jiang B, Parashar U, Nguyen T, Bines J, Patel MM. Childhood intussusception: a literature review. *PLoS ONE*. 2013;8:e68482.
2. Liu N, Yen C, Huang T, Cui P, Tate JE, Jiang B, et al. Incidence and epidemiology of intussusception among children under 2 years of age in Chenzhou and Kaifeng, China, 2009–2013. *Vaccine*. 2018;36:7862–7.
3. Gluckman S, Karpelowsky J, Webster AC, McGee RG. Management for intussusception in children. *Cochrane Database Syst Rev*. 2017;6:Cd006476.
4. Das MK, Arora NK, Mathai J, Sam CJ, G R. Profile and Epidemiology of Intussusception in Children under-two years of age: a prospective surveillance. *Indian J Pediatr*. 2021;88:1187–94.
5. Li XZ, Wang H, Song J, Liu Y, Lin YQ, Sun ZX. Ultrasonographic diagnosis of Intussusception in Children: a systematic review and Meta-analysis. *J Ultrasound Med*. 2021;40:1077–84.
6. Plut D, Phillips GS, Johnston PR, Lee EY. Practical imaging strategies for Intussusception in Children. *AJR Am J Roentgenol*. 2020;215:1449–63.
7. Kelley-Quon LI, Arthur LG, Williams RF, Goldin AB, St Peter SD, Beres AL, et al. Management of intussusception in children: a systematic review. *J Pediatr Surg*. 2021;56:587–96.
8. Sun Z, Song G, Lian D, Zhang Q, Dong L. Process management of Intussusception in Children: a retrospective analysis in China. *Pediatr Emerg Care*. 2022;38:321–5.
9. Niramis R, Watanatittan S, Kruatrachue A, Anuntkosol M, Buranakitjaroen V, Rattanasuwan T, et al. Management of recurrent intussusception: nonoperative or operative reduction? *J Pediatr Surg*. 2010;45:2175–80.
10. Fecteau A, Flageole H, Nguyen LT, Laberge JM, Shaw KS, Guttman FM. Recurrent intussusception: safe use of hydrostatic enema. *J Pediatr Surg*. 1996;31:859–61.

11. Gray MP, Li SH, Hoffmann RG, Gorelick MH. Recurrence rates after intussusception enema reduction: a meta-analysis. *Pediatrics*. 2014;134:110–9.
12. Esmaili-Dooki MR, Moslemi L, Hadipoor A, Osia S, Fatemi SA. Pediatric Intussusception in Northern Iran: comparison of recurrent with non-recurrent cases. *Iran J Pediatr*. 2016;26:e3898.
13. Guo WL, Hu ZC, Tan YL, Sheng M, Wang J. Risk factors for recurrent intussusception in children: a retrospective cohort study. *BMJ Open*. 2017;7:e018604.
14. Chen X, Chen Q, Wang X, Gao Z. Clinical characteristics of recurrent intussusception: a single-center retrospective study. *J Pediatr Surg*. 2021;56:1831–4.
15. Kim JH, Lee JS, Ryu JM, Lim KS, Kim WY. Risk factors for recurrent intussusception after fluoroscopy-guided Air Enema. *Pediatr Emerg Care*. 2018;34:484–7.
16. Xie X, Wu Y, Wang Q, Zhao Y, Xiang B. Risk factors for recurrence of intussusception in pediatric patients: a retrospective study. *J Pediatr Surg*. 2018;53:2307–11.
17. Vo A, Levin TL, Taragin B, Khine H. Management of Intussusception in the Pediatric Emergency Department: risk factors for recurrence. *Pediatr Emerg Care*. 2020;36:e185–e8.
18. Chen SC, Wang JD, Hsu HY, Leong MM, Tok TS, Chin YY. Epidemiology of childhood intussusception and determinants of recurrence and operation: analysis of national health insurance data between 1998 and 2007 in Taiwan. *Pediatr Neonatol*. 2010;51:285–91.
19. Hou J, Hou J, Die X, Sun J, Zhang M, Liu W, et al. Impact of overweight/obesity on clinical outcomes after reduction for intussusception in children. *Pediatr Surg Int*. 2021;37:887–95.
20. Zhang Y, Shao CC, Wei XL, Ni PJ, Guan H, Zhao C, et al. Ultrasound Findings to predict risk of recurrence in Pediatric Intussusception after Air Enema Reduction. *J Ultrasound Med*. 2022;41:1227–35.
21. Ye X, Tang R, Chen S, Lin Z, Zhu J. Risk factors for recurrent intussusception in children: a systematic review and Meta-analysis. *Front Pediatr*. 2019;7:145.
22. Tareen F, Ryan S, Avanzini S, Pena V, Mc Laughlin D, Puri P. Does the length of the history influence the outcome of pneumatic reduction of intussusception in children? *Pediatr Surg Int*. 2011;27:587–9.
23. Wang Z, He QM, Zhang H, Zhong W, Xiao WQ, Lu LW, et al. Intussusception patients older than 1 year tend to have early recurrence after pneumatic enema reduction. *Pediatr Surg Int*. 2015;31:855–8.
24. Chang D-H, Shin J, Rhee M-S, Park K-R, Cho B-K, Lee S-K et al. Vaginal microbiota profiles of native Korean women and associations with high-risk pregnancy. 2020.
25. Simanovsky N, Issachar O, Koplewitz B, Lev-Cohain N, Rekhman D, Hiller N. Early recurrence of ileocolic intussusception after successful air enema reduction: incidence and predisposing factors. *Emerg Radiol*. 2019;26:1–4.
26. Shen G, Zhang C, Li J, Zhang J, Liu Y, Guan Z, et al. Risk factors for short-term recurrent intussusception and reduction failure after ultrasound-guided saline enema. *Pediatr Surg Int*. 2018;34:1225–31.
27. Shen G, Liu H, Guan Z, Shang X, Li J, Zhang C, et al. Clinical features and factors leading to early recurrence of intussusception after saline reduction. *J Pak Med Assoc*. 2020;70:1727–30.
28. Zhu D, Xu X, Zhang M, Wang T, Zhu H. Significance of MCP-1 in predicting the short-term recurrence of primary intussusception in children: an observational study. *Med (Baltim)*. 2022;101:e30743.
29. Cho MJ, Nam CW, Choi SH, Hwang EH. Management of recurrent ileocolic intussusception. *J Pediatr Surg*. 2020;55:2150–3.
30. Adhikari S, Koirala DP, Pokhrel RP, Dahal GR, Kharel S, Neupane S. Risk factors for recurrent intussusception after successful reduction in pediatric patients in a tertiary care hospital of Nepal: a prospective study. *Ann Med Surg (Lond)*. 2022;76:103427.
31. Bartocci M, Fabrizi G, Valente I, Manzoni C, Specca S, Bonomo L. Intussusception in childhood: role of sonography on diagnosis and treatment. *J Ultrasound*. 2015;18:205–11.
32. Fiegel H, Gfroerer S, Rolle U. Systematic review shows that pathological lead points are important and frequent in intussusception and are not limited to infants. *Acta Paediatr*. 2016;105:1275–9.
33. Koumanidou C, Vakaki M, Pitsoulakis G, Kakavakis K, Mirilas P. Sonographic detection of lymph nodes in the intussusception of infants and young children: clinical evaluation and hydrostatic reduction. *AJR Am J Roentgenol*. 2002;178:445–50.

## Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.