

RESEARCH ARTICLE

Open Access



Diagnostic performance of individual characteristics and anthropometric measurements in detecting elevated serum alanine aminotransferase among children and adolescents

Yu-Lan Ou¹, Yue-Rong Lai¹, Chao-Nan Jiang¹, Jing Zhang² and Zan Ding^{2*}

Abstract

Background: Screening for elevated serum alanine aminotransferase (ALAT) can help identifying individuals at the risks of chronic and metabolic diseases, but blood collection is invasive and cannot be widely used for investigations. Considered as simple and inexpensive screening indices, individual characteristics and anthropometric measurements can be measured in a large crowd and may be important surrogate markers for ALAT levels. This study aimed to examine the diagnostic performance of individual characteristics and anthropometric parameters as predictive factors for discerning an elevated ALAT activity among Shenzhen children and adolescents.

Methods: A school-based screening study was performed from 9 high schools in Shenzhen during February 2017 and June 2018. Receiver operating characteristic curve was used to examine the diagnostic performance of each variable for detecting elevated ALAT.

Results: Altogether 7271 students aged 9–17 years were involved. The proportion of elevated ALAT greatly increased with increasing classification of BMI-z. By the sex-specific cut-offs for elevated ALAT (30 U/L boys; 19 U/L girls), BMI showed the highest area under the curve of 0.789 (95% CI 0.765–0.812) and followed by weight (0.779 [0.755–0.802]), BMI-z (0.747 [0.722–0.772]), height (0.622 [0.597–0.647]), and age (0.608 [0.584–0.632]), while height-z was not capable. With the cut-off of 67.8 kg for weight and 22.6 kg/m² for BMI, the accuracy to identify elevated ALAT was 87.1% for weight and 82.9% for BMI.

Conclusions: The presence of elevated ALAT was more common in overweight or obese children and adolescents. BMI and weight had the superiority of detecting elevated ALAT, followed by BMI-z, height, and age.

Keywords: Adolescent, Alanine aminotransferase, Anthropometric measurement, Body mass index, BMI-z

* Correspondence: dingzan_1990@163.com; dingzan@email.szu.edu.cn

²The Institute of Metabolic Diseases, Baoan Central Hospital of Shenzhen, The Fifth Affiliated Hospital of Shenzhen University, Shenzhen, Guangdong 518102, P.R. China

Full list of author information is available at the end of the article



© The Author(s). 2020 **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Background

The liver enzyme of serum alanine aminotransferase (ALAT or ALT) could be used clinically as a screening tool for the detection of probable nonalcoholic fatty liver disease (NAFLD), potential liver dysfunction, viral hepatitis infection, hepatocellular damage, and infectious mononucleosis [1–7]. An elevation of serum ALAT concentration is a key feature of NAFLD, and liver biopsy is the gold standard to confirm the diagnosis of NAFLD, characterized by pathological changes to liver structure and function such as fat accumulation, hepatocyte dysfunction, and fibrosis [7]. In addition, the elevation of ALAT has been recorded by published evidence to be strongly linked to, or to predict, the developing of type 2 diabetes mellitus, insulin resistance, and cardiovascular diseases (e.g. atherogenesis, congestive heart failure and coronary heart disease) [5, 6, 8–14].

Screening for the concentration of serum ALAT and verifying an elevated ALAT activity could assist in determining individuals at risk for the above chronic conditions and reducing the possibility of future liver diseases [15, 16]. However, the measurement of ALAT levels requires blood collection, which is invasive and could be impractical in large-scale epidemiological investigations because of subject burden and cost. In contrast, individual characteristic (e.g. age and gender) and measurements of anthropometric indices (e.g. height, weight, body mass index [BMI], and waist circumference [WC]) are relatively simple, cheap, quick, and non-invasive, and therefore can be applied to a great number of people and easily conducted during common health examinations, especially for non-adults [15, 17, 18]. If these indices showed a close correlation with serum ALAT levels or had sufficient ability to detect an ALAT elevation, they could be a useful surrogate for ALAT levels.

Two multi-center, population-based epidemiological researches conducted in the mainland of China have consistently demonstrated that the higher values of BMI, hip circumference, WC, waist-to-hip ratio (WHpR), and waist-to-height ratio (WHtR) were able to predict an elevated ALAT activity [17, 19]. Recently, the nontraditional index of BMI z-score has also been put forward as another important anthropometric indicator of ALAT elevation [7, 20]. However, whether the potential of the z-score of height, a novel body index, for predicting elevated ALAT was comparable or superior to the abovementioned parameters was still unclear up to now. In addition, although most previous studies on the relations of anthropometric measurements and individual characteristics with ALAT elevation were conducted among adults [17, 21–23], few investigations focused on the performance predication of those predictors for diagnosing elevated ALAT among children and adolescents [15, 20, 24, 25].

Based on a school-based screening, the current study mainly aimed to examine the diagnostic performance of individual characteristics (i.e. age and gender) and several anthropometric parameters (i.e. height, weight, BMI, BMI-z, and height-z) as predictive factors of an elevated ALAT activity in Shenzhen children and adolescents, and to determine the optimal cut-off points for these parameters that would identify a person with an elevation of serum ALAT level. Assessing individual indicators is crucial for early detection and prevention of an activity of elevated ALAT or the subsequent development of NAFLD among children and adolescents.

Methods

Study setting and population

Shenzhen is one of the 4 first-tier cities in mainland China and a major financial centre and the world's largest manufacturing base [26]. The population and economy have grown rapidly, with a total gross domestic product of 255 billion USD in 2015 [27].

Data collections and measures

Involved freshmen from 9 junior and senior high schools of Bao'an District, a cross-sectional study organized with the assistance of the Bao'an Government of Shenzhen was officially carried out [28]. Data collection activities were completed during February 2017 and June 2018. This school-based study was approved by the Institutional Review Board of Baoan Central Hospital of Shenzhen, and written informed consents for each student and their parents were obtained.

Individual characteristic data such as gender and date of birth were taken from each participant, and anthropometric parameters of weight and standing-height were respectively measured to the nearest 0.1 kg and 0.5 cm by trained physicians using a standard weighing machine. At each high school, blood samples were drawn in the morning after participants fasting for at least 10 h by trained nurses. ALAT was measured with the fully automatic biochemical analyzer (Model AU5821, Tokyo, Japan). Finally, a total of 7271 students who had complete anthropometric and clinical data participated.

For each participant, age with 1 decimal was transformed based on the date of birth and BMI (kg/m^2) was calculated in weight divided by the square of height. Based on an international norm from the World Health Organization growth reference (i.e. the 2007 WHO reference) for school-aged children and adolescents aged 5–19 years, anthropometric indicators of z-score values of height-for-age (height-z) and BMI-for-age (BMI-z) for both boys and girls were calculated by the R macro, provided on the WHO website (<https://www.who.int/growthref/en/>) [29, 30]. Moreover, according to the z-score of BMI-for-age, we defined BMI-z > 2SD as obesity, 1SD <

BMI- $z \leq 2SD$ as overweight, $-2SD \leq \text{BMI-}z \leq 1SD$ as normal-weight, and $\text{BMI-}z < -2SD$ as underweight; where SD was standard deviation of the BMI z-scores.

Definition of elevated serum ALAT levels

The thresholds of elevated ALAT needed to be determined consistent with previously reported levels, as there was no consensus on what level constituted an elevated serum ALAT concentration for children and adolescents. For the most commonly use of definitions among children and adolescents, an activity of elevated serum ALAT was defined as > 30 U/L for boys and > 19 U/L for girls (diagnostic criterion I) [5, 6, 11, 31]. Main analysis was undertaken using these sex-specific cut-offs, and a secondary analysis for the sake of comparison was also reported, defining with higher thresholds (> 40 U/L for both boys and girls; diagnostic criterion II) that have been proposed in previous observational literatures [10, 16, 32] to explore whether higher thresholds led to the same or different results.

Statistical analyses

The normal distribution of each continuous measurement was determined by the one-sample Kolmogorov-Smirnov test. The non-normally distributed measurements were described as median with inter-quartile range (25th percentile–75th percentile; IQR); significance for differences between elevated and normal serum ALAT was evaluated by the Mann–Whitney U test. Stratified by the classification of BMI z-score, the crude prevalence along with 95% confidence interval (CI) of elevated ALAT was quantitatively estimated, and the trend of the proportion of elevated ALAT based on BMI- z was carried out with chi-square test. The Spearman's rank correlation analyses among the skewed parameters were performed.

Extensively used in clinical epidemiology for evaluation of diagnostic ability of biomarkers (e.g. serum markers) or a diagnostic test with dichotomous outcome (i.e. positive or negative result) in classification of the diseased from healthy population, the receiver operating characteristic (ROC) curve, defined as a plot of the sensitivity of a test as y-axis versus 1-specificity as x-axis, is an effective method and graphical technique to describe the accuracy of a prediction model or diagnostic test [33, 34]. Sensitivity and specificity, the basic measures of accuracy of a diagnostic test, vary with different cut-off thresholds, and sensitivity is inversely related to specificity [18, 34]. Positive predictive value (PPV) is defined as the probability of disease for positive test results, and negative predictive value (NPV) is defined as the probability of being healthy for negative test results [33].

In the current study, we utilized the ROC curve analysis to find out the optimal cut-off points and to

examine the diagnostic performance of each measurement as indicators of ALAT elevation. An anthropometric parameter value with the highest Youden index was chosen as the optimal cut-off point [18]. The observed agreement and the area under the curve (AUC) were also determined. Statistical analyses were processed in R 3.5.1 (<http://www.R-project.org>) and SPSS for Windows 16.0, with two-tailed P -value < 0.05 considered statistically significant.

Results

Characteristics of participants

In the current investigation, a total of 7271 students aged 9–17 years were recruited for the operative statistical analysis, involving 4014 (55.2%) boys. Characteristics of study subjects with and without elevated ALAT activity are listed in Table 1. The indices of age, weight, height, BMI, height- z , and BMI- z were non-normal distribution (all $P < 0.001$). Overall, the median (IQR) level was 14.7 (12.4–15.7) years for age, 53.0 (46.7–61.0) kg for weight, 1.65 (1.59–1.72) m for height, 19.31 (17.63–21.66) kg/m^2 for BMI, 0.59 (–0.18 to 1.41) for height- z , and –0.01 (–0.77 to 0.83) for BMI- z . Each anthropometric index but not height- z was significantly greater among students with an elevated ALAT activity than those with a normal ALAT level, regardless of the use of definitions (most $P < 0.001$).

Crude prevalence of elevated ALAT by the BMI- z classification

On the basis of ALAT activity thresholds of diagnostic criterion I, an elevated ALAT activity was present in 2.48% (6/242) of underweight, 3.79% (208/5493) of normal-weight, 12.58% (137/1089) of overweight, and 36.91% (165/447) of obese children and adolescents; based on the criterion II, the proportion of elevated ALAT was 0.83% (2/242), 0.89% (49/5493), 4.96% (54/1089), and 20.58% (92/447) for the corresponding subgroups (Table S1). Regardless of the use of diagnostic criteria, the abnormality prevalence rate of ALAT greatly increased with increasing classification of BMI z-score and peaked in the obesity group for all participants, boys, and girls (all $P_{trend} < 0.001$). Overweight or obese students were much more likely to obtain increased ALAT concentrations than normal-weight or underweight students.

Univariate ROC curve analyses

Figure 1 illustrates the diagnostic performance of individual indices for identifying elevated ALAT by ROC curves, and Table 2 presents the information about the AUCs of the curves among Shenzhen children and adolescents. The anthropometric parameter of BMI showed the highest AUC (95% CI) for elevated ALAT (0.789

Table 1 Baseline descriptive statistics for individual measurements stratified by gender among Shenzhen children and adolescents (aged 9–17 years), separately based on the diagnostic criteria I and II for elevated ALAT

Group	Total students	Criterion I (> 30 U/L for boys and > 19 U/L for girls)			Criterion II (> 40 U/L for boys and girls)		
		Normal ALAT	Elevated ALAT	P-value*	Normal ALAT	Elevated ALAT	P-value*
Overall (n = 7271)							
Age, years	14.7 (12.4–15.7)	14.7 (12.3–15.7)	15.3 (13.8–16.1)	< 0.001	14.7 (12.3–15.7)	15.3 (14.3–16.1)	< 0.001
Height, m	1.65 (1.59–1.72)	1.64 (1.59–1.71)	1.69 (1.63–1.75)	< 0.001	1.65 (1.59–1.72)	1.73 (1.65–1.78)	< 0.001
Weight, kg	53.0 (46.7–61.0)	52.2 (46.2–59.8)	70.0 (56.1–83.6)	< 0.001	52.6 (46.5–60.2)	77.7 (63.1–90.1)	< 0.001
BMI, kg/m ²	19.31 (17.63–21.66)	19.15 (17.55–21.23)	24.51 (20.38–27.95)	< 0.001	19.25 (17.59–21.45)	26.39 (22.40–30.22)	< 0.001
Height-z	0.59 (–0.18–1.41)	0.58 (–0.18–1.41)	0.71 (–0.12–1.47)	0.104	0.59 (–0.18–1.41)	0.65 (–0.13–1.48)	0.526
BMI-z	–0.01 (–0.77–0.83)	–0.06 (–0.81–0.74)	1.40 (0.16–2.27)	< 0.001	–0.04 (–0.79–0.79)	1.89 (0.85–2.54)	< 0.001
Boys (n = 4014)							
Age, years	14.7 (12.4–15.7)	14.7 (12.3–15.7)	15.4 (14.4–16.1)	< 0.001	14.7 (12.4–15.7)	15.4 (14.6–16.2)	< 0.001
Height, m	1.69 (1.61–1.75)	1.69 (1.61–1.75)	1.74 (1.68–1.78)	< 0.001	1.69 (1.61–1.75)	1.74 (1.68–1.79)	< 0.001
Weight, kg	55.4 (48.1–64.7)	54.7 (47.9–63.0)	79.0 (66.9–88.6)	< 0.001	55.0 (48.0–63.7)	81.4 (69.5–92.4)	< 0.001
BMI, kg/m ²	19.37 (17.64–22.04)	19.16 (17.55–21.47)	26.42 (22.98–30.06)	< 0.001	19.26 (17.59–21.70)	27.20 (23.92–30.61)	< 0.001
Height-z	0.65 (–0.16–1.55)	0.65 (–0.16–1.55)	0.66 (–0.16–1.42)	0.086	0.65 (–0.16–1.55)	0.66 (–0.12–1.50)	0.857
BMI-z	0.12 (–0.74–1.06)	0.04 (–0.78–0.90)	1.98 (0.99–2.55)	< 0.001	0.07 (–0.77–0.95)	2.14 (1.27–2.67)	< 0.001
Girls (n = 3257)							
Age, years	14.7 (12.3–15.7)	14.7 (12.3–15.7)	15.0 (13.3–16.0)	< 0.001	14.7 (12.3–15.7)	14.6 (13.3–15.4)	0.557
Height, m	1.62 (1.57–1.67)	1.61 (1.57–1.66)	1.64 (1.58–1.71)	< 0.001	1.62 (1.57–1.66)	1.62 (1.58–1.69)	0.270
Weight, kg	50.6 (45.2–57.0)	50.1 (45.0–56.0)	58.5 (50.6–71.1)	< 0.001	50.5 (45.2–56.8)	59.0 (48.7–75.6)	< 0.001
BMI, kg/m ²	19.26 (17.60–21.26)	19.13 (17.54–20.98)	21.60 (19.05–25.59)	< 0.001	19.23 (17.60–21.19)	22.01 (18.86–27.12)	< 0.001
Height-z	0.51 (–0.20–1.32)	0.50 (–0.20–1.31)	0.82 (–0.09–1.52)	0.006	0.51 (–0.20–1.32)	0.40 (–0.20–1.37)	0.857
BMI-z	–0.13 (–0.82–0.61)	–0.16 (–0.84–0.54)	0.64 (–0.39–1.60)	< 0.001	–0.14 (–0.82–0.60)	0.73 (–0.53–1.80)	< 0.001

Data are expressed as median (interquartile range [IQR]) for the measurements with skewed distributions

* Comparing elevated ALAT to normal ALAT by the Mann–Whitney U test

[0.765–0.812] by criterion I; 0.850 [0.818–0.882] by criterion II), and followed by weight (0.779 [0.755–0.802]; 0.850 [0.817–0.882]) and the z-score of BMI (0.747 [0.722–0.772]; 0.822 [0.787–0.858]). Compared to the above 3 obesity indices, age and height had a poorer diagnostic efficiency in detecting abnormal ALAT. The discriminatory power of height z-score in the prediction of elevated ALAT was null (both $P > 0.1$), and gender also had no ability to evaluate elevated ALAT by the diagnostic criterion I ($P = 0.435$). In the stratification analyses by gender, similar outcomes were provided in Table S2.


Table 3 elucidates the optimal cut-off points and the probability of meaningful and significant parameters in diagnosing abnormal ALAT, based on ROC analyses. The suggested cut-off points of were 67.8 & 68.9 kg for weight and 22.6 & 22.3 kg/m² for BMI, separately by the criteria I and II. Based on the cut-offs, the parameter of weight achieved the highest ability to correctly classify subjects as elevated ALAT, with an accuracy of 87.1% by criterion I and 88.9% by criterion II. However, BMI took control of a larger Youden index (46.1%) than other parameters with criterion I. BMI and weight had an

approximate diagnostic performance in diagnosing elevated ALAT.

Correlation analyses and combined individual measurements for detecting elevated ALAT in multivariate analysis

Table S3 displays the Spearman’s correlation coefficients between individual indices and ALAT levels. BMI had a very high positive correlation with BMI z-score ($r = 0.891$) and weight ($r = 0.829$), suggesting a multicollinearity among each obesity parameter. Weight was also quite strongly and positively correlated with height ($r = 0.673$) and BMI-z ($r = 0.657$).

Table 4 shows the capability of combined individual indices to diagnose the presence of elevated ALAT in multivariate models. The combined indices of “age + gender+height + weight” showed the strongest capability to indicate the levels of ALAT, with an AUC (95% CI) of 0.816 (0.795–0.836) for criterion I and 0.858 (0.826–0.891) for criterion II. In particular, a very slight increase of 0.037 (0.816 vs 0.779) for criterion I and 0.008 (0.858 vs 0.850) for criterion II in accuracy was demonstrated, as compared with the univariate weight model. The



multivariate combined BMI model also fitted well with an analogous good accuracy (AUC = 0.813 for criterion I and 0.858 for II). Compared to the univariate models of obesity indices (i.e. weight, BMI, or BMI-z), the combined models did not markedly improve the prediction of elevated ALAT for children and adolescents.

Discussion

Primary focused on the discriminative accuracy of the indices, individual characteristic and anthropometric data for Shenzhen children and adolescents aged 9–17 years were used to diagnose elevated ALAT, and we confirmed the belief that individual indices were important predictors of abnormal ALAT. An elevated ALAT

activity was more common in overweight or obese students than those with a normal or relatively low BMI z-score. BMI and weight showed an approximate diagnostic performance for predicting the presence of elevated ALAT, followed by the z-score of BMI, height, and age.

Consistent findings that the proportion of elevated ALAT greatly increased with increasing degree of obesity (e.g. the percentile or z-score of BMI) and higher proportion among overweight or obese adolescents than normal-weight ones were shown in previous studies and our research, regardless of the use of diagnostic criteria for an elevated ALAT activity [10, 16, 21, 35]. Based on the criterion II (40 U/L for boys and girls), the percentage of elevated ALAT among Shenzhen children and

Table 2 The AUC (95% CI) of the ROC curve to judge the discrimination ability of various individual measurements for detecting elevated ALAT (separately by the criteria I and II) in univariate analyses

Variable	Criterion I (> 30 U/L for boys and > 19 U/L for girls)				Criterion II (> 40 U/L for boys and girls)			
	AUC	95% CI	SE	P*	AUC	95% CI	SE	P*
Age	0.608	0.584–0.632	0.012	< 0.001	0.613	0.577–0.649	0.018	< 0.001
Gender	0.490	0.464–0.516	0.013	0.435	0.628	0.593–0.664	0.018	< 0.001
Height	0.622	0.597–0.647	0.013	< 0.001	0.695	0.658–0.732	0.019	< 0.001
Weight	0.779	0.755–0.802	0.012	< 0.001	0.850	0.817–0.882	0.016	< 0.001
BMI	0.789	0.765–0.812	0.012	< 0.001	0.850	0.818–0.882	0.016	< 0.001
Height-z	0.522	0.496–0.547	0.013	0.102	0.514	0.473–0.554	0.021	0.517
BMI-z	0.747	0.722–0.772	0.013	< 0.001	0.822	0.787–0.858	0.018	< 0.001

AUC Area under curve, ALAT Alanine aminotransferase, CI Confidence interval, ROC Receiver operating characteristic, SE Standard error

* Null hypothesis: true area = 0.5

Table 3 The optimal cut-off points and the probability of significant individual parameters for identifying elevated ALAT, separately based on the diagnostic criteria I and II

Parameter	Criterion I (> 30 U/L for boys and > 19 U/L for girls)							Criterion II (> 40 U/L for boys and girls)						
	Cut-off	Accuracy	Sen	Spe	Youden index	PPV	NPV	Cut-off	Accuracy	Sen	Spe	Youden index	PPV	NPV
Age	13.7 years	44.0	76.2 ^a	41.5	17.7	9.0	95.8	13.7 years	41.8	78.7 ^a	40.8	19.5	3.6	98.6
Height	164.0 cm	49.0	72.3	47.3	19.5	9.5	95.7	169.5 cm	67.3	61.9	67.5	29.4	5.0	98.5
Weight	67.8 kg	87.1 ^a	55.0	89.5 ^a	44.6	28.6 ^a	96.3	68.9 kg	88.9 ^a	69.5	89.5 ^a	59.0 ^a	15.6 ^a	99.1
BMI	22.6 kg/m ²	82.9	61.6	84.5	46.1 ^a	23.3	96.6 ^a	22.3 kg/m ²	80.8	77.2	80.9	58.0	10.1	99.2 ^a
BMI-z	0.95	79.1	60.1	80.5	40.6	19.1	96.4	1.25	84.3	70.1	84.7	54.7	11.3	99.0

The optimal cut-off points were based on the maximum Youden index for each parameter; the probability of parameters for identifying elevated ALAT is expressed as %

Sen Sensitivity, Spe Specificity, PPV Positive predictive value, NPV Negative predictive value

^a The best performance for each conventional diagnostic test index

adolescents of 9–17 years was 0.89% for normal-weight, 4.96% for overweight, and 20.58% for obese participants. According to the same thresholds, elevated ALAT levels were observed in nearly 6.6% of the Mexican youths (9.8% of boys and 3.8% of girls), and 2.7% of normal-BMI, 14.2% of overweight, and 28.9% of the obese children and adolescents, by using the baseline data from 1262 participants of 8–19 years in the Mexican Health Worker Cohort Study [16]. Mexican youths had a slightly higher crude proportion of elevated ALAT than Shenzhen youths, without adjusting for age and gender based on the distribution of the world population. Likewise, using a sample of 1591 youths from the 2008–2009 Korea National Health and Nutrition Examination Survey, another study conducted by Seung Park and cooperators also found a close prevalence of elevated ALAT (> 33 U/L for boys and > 25 U/L for girls) among Korean youths of 12–18 years — 5.9% (95%CI 4.9%–7.2%) for the overall study population, 15.7% (11.3%–21.5%) for overweight adolescents (85th ≤ BMI < 95th percentile), and 34.9 (25.6%–45.4%) for subjects with a BMI ≥ 95th percentile [35]. The odds ratios (ORs) for elevated ALAT also sharply increased with the greater levels of obesity, with an OR (95% CI) of 7.23 (4.33–12.10) for overweight and 23.62 (12.98–42.98) for obese adolescents, comparing to normal-weight adolescents (BMI < 85th percentile) in an unadjusted analysis [35].

Screening for school-students, it is clearly more convenient to collect anthropometric measures than biochemical measures, but the abilities of several individual characteristics and anthropometric indices to correctly predict elevated serum ALAT in children and adolescents are questionable and need to be assessed. Determined by the univariate ROC curves, our results showed that the anthropometric parameter of BMI had the best superiority of discerning the presence of elevated ALAT (AUC = 0.789 for criterion I and 0.850 for II), weight and BMI-z displayed the second and third highest detection accuracy (from 0.747 to 0.850), and followed by height, age, and gender (from 0.490 to 0.695). As such, we believed that the single variable of height, age, or gender was not a reliable surrogate measure of an activity of elevated ALAT among Shenzhen students of 9–17 years. Further, age, gender, and height were considered as covariates together with each predictor of obesity indices (i.e. weight, BMI, or BMI-z) to predict elevated ALAT in the subsequent multivariate model analyses, although the diagnostic accuracy of the combined models did not markedly improve. On the other hand, the current study firstly investigated the association of height z-score with serum ALAT and estimated the usefulness of height-z as a predictive index of elevated ALAT among children and adolescents, showing a null discriminative power of height z-score for predicting an elevated ALAT activity.

Table 4 The AUC (95% CI) of the ROC curve to express the diagnostic performance of combined individual measurements for detecting elevated ALAT (separately by the diagnostic criteria I and II) in multivariate analyses

Combined measurements	Criterion I (> 30 U/L for boys and > 19 U/L for girls)				Criterion II (> 40 U/L for boys and girls)			
	AUC	95% CI	SE	P*	AUC	95% CI	SE	P*
Age + gender+height	0.643	0.619–0.668	0.012	< 0.001	0.716	0.679–0.753	0.019	< 0.001
Age + gender+height + weight	0.816	0.795–0.836	0.011	< 0.001	0.858	0.826–0.891	0.016	< 0.001
Age + gender+height + BMI	0.813	0.792–0.834	0.011	< 0.001	0.858	0.826–0.890	0.016	< 0.001
Age + gender+height + BMI-z	0.807	0.785–0.829	0.011	< 0.001	0.856	0.823–0.889	0.017	< 0.001

AUC Area under curve, ALAT Alanine aminotransferase, CI Confidence interval, ROC Receiver operating characteristic, SE Standard error

* Null hypothesis: true area = 0.5

For BMI, several studies had pointed out that BMI was able to determine the presence of elevated ALAT among adults, although the accuracy was not very high in absolute term — the AUC (95% CI) was 0.658 (0.633–0.683) for men and 0.651 (0.616–0.685) for women in the rural areas of China [17] and 0.64 (0.60–0.68) for Italian general population [22]. Focused on adolescents, a representative study with a sample of 454 youths of 11–17 years from 2 northern Italian cities also found a similar diagnostic performance of dichotomized BMI for elevated ALAT, with an AUC of 0.64 (0.50–0.77) [20]. Compared to the dichotomized BMI model, the Italian adolescents study also indicated a more accurate univariate model of BMI-*z* (AUC = 0.71 [0.59–0.81]), and the predictive ability was increased substantially by considering gender together with BMI-*z* (AUC = 0.80 [0.71–0.89]) [20]. In addition, based on the data from the National Health and Nutrition Examination Survey during 1999 to 2014, an United States study consisted of 5019 adolescents of 12–19 years suggested a significant correlation between BMI *z*-score and serum ALAT levels ($r = 0.29$, $P < 0.0001$) [7]. Significant positive associations of ALAT with obesity indices (e.g. BMI, body fat percentage, truncal fat mass, total fat mass, WC, and WHtR) were further confirmed in Korean male adolescents [24].

Several potential limitations of our research should be recognized. First, participants were only from a single city of China, and it might be therefore difficult to generalize these findings to other populations. Second, our cross-sectional study omitted to measure some important individual anthropometric parameters — hip circumference, WC, neck circumference, and the subsequent indicators of WHpR, WHtR, and A Body Shape Index, which might achieve a better diagnostic performance and to be more sensitive predictive indicators for diagnosing an elevated ALAT activity among the indigenous adolescents. For predicting the ALAT levels, WHtR and to some extent BMI were congruously shown to be the best body indices in some Asians, as compares to WC, hip circumference, WHpR, and the useless index of A Body Shape Index [15, 17, 19]. Third, our study did not take into account some potential confounders of ALAT elevation for non-adults such as physical exercise, drugs, and dietary choline deficiency [1]. Another limitation was our inability to consider the potential confounders of ethyl alcohol intake and hepatitis B virus (HBV) and C virus (HCV) infections, which were well-known risk factors for increases in ALAT [7, 15, 20, 22]. However, alcohol consumption and HBV and HCV infections were less prevalent in school-students of China [36, 37], and we hypothesized that the obesity indices could be even more important predictors of elevated ALAT among the children and adolescents of Shenzhen, and the above

potential confounders were not likely to have a substantial impact on current results.

Conclusions

An elevated ALAT activity was more frequent in overweight or obese students than normal-weight subjects. Anthropometric obesity indices of BMI, weight, and BMI-*z* achieved a very high diagnostic performance for predicting elevated ALAT among Shenzhen children and adolescents and followed by height and age, while height-*z* was not capable.

Supplementary information

Supplementary information accompanies this paper at <https://doi.org/10.1186/s12887-020-02033-9>.

Additional file 1: Table S1. Crude prevalence of elevated ALAT stratified by the classification of BMI *z*-score among children and adolescents of Shenzhen, separately based on the diagnostic criteria I and II. **Table S2.** The AUC (95% CI) of the ROC curve stratified by gender in univariate analyses, separately based on the diagnostic criteria I and II for elevated ALAT. **Table S3.** Spearman's rank correlation coefficients between individual measurements and ALAT.

Abbreviations

ALAT: Alanine aminotransferase; AUC: Area under the curve; BMI: Body mass index; CI: Confidence interval; HBV: Hepatitis B virus; HCV: Hepatitis C virus; NPV: Negative predictive value; NAFLD: Nonalcoholic fatty liver disease; PPV: Positive predictive value; ROC: Receiver operating characteristic; SD: Standard deviation; WC: Waist circumference; WHpR: Waist-to-hip ratio; WHtR: Waist-to-height ratio

Acknowledgements

Not applicable.

Authors' contributions

Y.L.O. and Z.D. conceived and designed the experiments. Y.L.O., Y.R.L., C.N.J., and J.Z. collected and cleaned the data. Y.L.O. drafted the manuscript. Z.D. guided statistical analysis, revised the manuscript, and interpreted the results. All authors read and approved the final manuscript.

Funding

The study was supported by the Metabolic Disease in Prevention and Treatment Program of Occupational Groups and Adolescents (2019–2020), under the Bao'an Government of Shenzhen.

Availability of data and materials

Readers can get the datasets and materials of the current study by contacting the corresponding author of Zan Ding (dingzan_1990@163.com) for a reasonable request.

Ethics approval and consent to participate

This school-based study was approved by the Institutional Review Board of Baoan Central Hospital of Shenzhen, and written informed consents for each student and their parents were obtained.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Author details

¹Department of Gynecology, Sun Yat-sen University Cancer Center, Guangzhou 510060, Guangdong, China. ²The Institute of Metabolic Diseases,

Baoan Central Hospital of Shenzhen, The Fifth Affiliated Hospital of Shenzhen University, Shenzhen, Guangdong 518102, P.R. China.

Received: 14 January 2020 Accepted: 12 March 2020

Published online: 20 March 2020

References

- Loprinzi PD. Physical activity with alanine aminotransferase and gamma-glutamyltransferase among congestive heart failure patients. *Int J Cardiol.* 2016;212:61–2.
- Khalili M, Shuhart MC, Lombardero M, Feld JJ, Kleiner DE, Chung RT, et al. Relationship between metabolic syndrome, alanine aminotransferase levels, and liver disease severity in a multiethnic north American cohort with chronic hepatitis B. *Diabetes Care.* 2018;41:1251–9.
- Ghany MG, Feld JJ, Chang KM, Chan HL, Lok AS, Visvanathan K, et al. Serum alanine aminotransferase flares in chronic hepatitis B infection: the good and the bad. *Lancet Gastroenterol Hepatol.* 2020. [https://doi.org/10.1016/S2468-1253\(19\)30344-9](https://doi.org/10.1016/S2468-1253(19)30344-9).
- Liu Z, Que S, Xu J, Peng T. Alanine aminotransferase-old biomarker and new concept: a review. *Int J Med Sci.* 2014;11:925–35.
- Di Bonito P, Sanguigno E, Di Fraia T, Forziato C, Boccia G, Saitta F, et al. Association of elevated serum alanine aminotransferase with metabolic factors in obese children: sex-related analysis. *Metabolism.* 2009;58:368–72.
- Park HK, Hwang JS, Moon JS, Lee JA, Kim DH, Lim JS. Healthy range of serum alanine aminotransferase and its predictive power for cardiovascular risk in children and adolescents. *J Pediatr Gastroenterol Nutr.* 2013;56:686–91.
- Fermin CR, Lee AM, Filipp SL, Gurka MJ, DeBoer MD. Serum alanine aminotransferase trends and their relationship with obesity and metabolic syndrome in United States adolescents, 1999–2014. *Metab Syndr Relat Disord.* 2017;15:276–82.
- Song BM, Kim HC, Kim DJ, Ahn SV, Kim KM, Lee JM, et al. Aminotransferase levels, body mass index, and the risk of diabetes: a prospective cohort study. *Ann Epidemiol.* 2018;28:675–80.e6.
- Siddiqui MS, Sterling RK, Luketic VA, Puri P, Stravitz RT, Bouneva I, et al. Association between high-normal levels of alanine aminotransferase and risk factors for atherogenesis. *Gastroenterology.* 2013;145:1271–9.e1–3.
- Elizondo-Montemayor L, Ugalde-Casas PA, Lam-Franco L, Bustamante-Careaga H, Serrano-Gonzalez M, Gutierrez NG, et al. Association of ALT and the metabolic syndrome among Mexican children. *Obes Res Clin Pract.* 2014;8:e79–87.
- Samani SG, Keshladi R, Adibi A, Noori H, Moeini M. Association of serum alanine aminotransferase levels with cardiometabolic risk factors in normal-weight and overweight children. *Iran J Pediatr.* 2011;21:287–93.
- Williams KH, Burns K, Twigg SM. Differing clinical phenotype for higher alanine-aminotransferase (ALT) compared with high-risk NAFLD fibrosis score in type 2 diabetes mellitus. *J Diabetes Complicat.* 2018;32:321–4.
- Kong AP, Choi KC, Cockram CS, Ho CS, Chan MH, Ozaki R, et al. Independent associations of alanine aminotransferase (ALT) levels with cardiovascular risk factor clustering in Chinese adolescents. *J Hepatol.* 2008;49:115–22.
- Gomez-Samano MA, Cuevas-Ramos D, Mehta R, Brau-Figueroa H, Meza-Arana CE, Gullas-Herrero A. Association of Alanine Aminotransferase Levels (ALT) with the Hepatic Insulin Resistance Index (HIRI): a cross-sectional study. *BMC Endocr Disord.* 2012;12:16.
- Ochiai H, Shirasawa T, Nishimura R, Nanri H, Ohtsu T, Hoshino H, et al. Waist-to-height ratio is more closely associated with alanine aminotransferase levels than body mass index and waist circumference among population-based children: a cross-sectional study in Japan. *BMC Pediatr.* 2015;15:59.
- Purcell M, Flores YN, Zhang ZF, Denova-Gutierrez E, Salmeron J. Prevalence and predictors of alanine aminotransferase elevation among normal weight, overweight and obese youth in Mexico. *J Dig Dis.* 2013;14:491–9.
- Chen S, Guo X, Yu S, Zhou Y, Li Z, Sun Y. Anthropometric indices in adults: which is the best indicator to identify alanine aminotransferase levels? *Int J Environ Res Public Health.* 2016;13:226.
- Asif M, Aslam M, Altaf S. Evaluation of anthropometric parameters of central obesity in Pakistani children aged 5–12 years, using receiver operating characteristic (ROC) analysis. *J Pediatr Endocrinol Metab.* 2018. <https://doi.org/10.1515/jpem-2018-0193>.
- Wu XY, Hu CL, Wan YH, Su PY, Xing C, Qi XY, et al. Higher waist-to-height ratio and waist circumference are predictive of metabolic syndrome and elevated serum alanine aminotransferase in adolescents and young adults in mainland China. *Public Health.* 2012;126:135–42.
- Bedogni G, Miglioli L, Masutti F, Castiglione A, Tiribelli C, Bellentani S. Accuracy of body mass index in detecting an elevated alanine aminotransferase level in adolescents. *Ann Hum Biol.* 2004;31:570–7.
- Adams LA, Knuiam MW, Divitini ML, Olynyk JK. Body mass index is a stronger predictor of alanine aminotransferase levels than alcohol consumption. *J Gastroenterol Hepatol.* 2008;23:1089–93.
- Bedogni G, Miglioli L, Battistini N, Masutti F, Tiribelli C, Bellentani S. Body mass index is a good predictor of an elevated alanine transaminase level in the general population: hints from the Dionysos study. *Dig Liver Dis.* 2003;35:648–52 (in eng).
- Kim J, Jo I. Relationship between body mass index and alanine aminotransferase concentration in non-diabetic Korean adults. *Eur J Clin Nutr.* 2010;64:169–75.
- Ahn MB, Bae WR, Han KD, Cho WK, Cho KS, Park SH, et al. Association between serum alanine aminotransferase level and obesity indices in Korean adolescents. *Korean J Pediatr.* 2015;58:165–71.
- Bussler S, Vogel M, Pietzner D, et al. New pediatric percentiles of liver enzyme serum levels (alanine aminotransferase, aspartate aminotransferase, γ -glutamyltransferase): effects of age, sex, body mass index, and pubertal stage. *Hepatology (Baltimore, Md).* 2018;68:1319–30.
- Zhan ZY, Yu YM, Qian J, Song YF, Chen PY, Ou CQ. Effects of ambient temperature on ambulance emergency call-outs in the subtropical city of Shenzhen, China. *PLoS One.* 2018;13:e0207187.
- Guo Y, Ma Y, Ji J, Liu N, Zhou G, Fang D, et al. The relationship between extreme temperature and emergency incidences: a time series analysis in Shenzhen, China. *Environ Sci Pollut Res Int.* 2018;25:36239–55.
- Ding Z, Zhang J, Deng C-Y, You Y-B, Zhou H. Association of body mass index with serum alanine aminotransferase in Chinese adolescents: a school-based cross-sectional study. *Gastroenterol Rep.* 2019. <https://doi.org/10.1093/gastro/goz039>.
- de Onis M, Onyango AW, Borghi E, Siyam A, Nishida C, Siekmann J. Development of a WHO growth reference for school-aged children and adolescents. *Bull World Health Organ.* 2007;85:660–7.
- Zong XN, Li H. Construction of a new growth references for China based on urban Chinese children: comparison with the WHO growth standards. *PLoS One.* 2013;8:e59569.
- Prati D, Taioli E, Zanella A, Della Torre E, Butelli S, Del Vecchio E, et al. Updated definitions of healthy ranges for serum alanine aminotransferase levels. *Ann Intern Med.* 2002;137:1–10.
- Ramirez-Lopez G, Moran-Villota S, Mendoza-Carrera F, Portilla-de Buen E, Valles-Sanchez V, Castro-Martinez XH, et al. Metabolic and genetic markers' associations with elevated levels of alanine aminotransferase in adolescents. *J Pediatr Endocrinol Metab.* 2018;31:407–14.
- Hajian-Tilaki K. Receiver operating characteristic (ROC) curve analysis for medical diagnostic test evaluation. *Caspian J Intern Med.* 2013;4:627–35.
- Obuchowski NA, Bullen JA. Receiver operating characteristic (ROC) curves: review of methods with applications in diagnostic medicine. *Phys Med Biol.* 2018;63:07tr1.
- Park SH, Heo NY, Park JH, Kim TO, Yang SY, Moon YS, et al. Obesity, insulin resistance, and the risk of an elevated alanine aminotransferase activity in the Korean adolescent population. *J Pediatr Endocrinol Metab.* 2012;25:945–9.
- Guo L, Deng J, He Y, Deng X, Huang J, Huang G, et al. Alcohol use and alcohol-related problems among adolescents in China: a large-scale cross-sectional study. *Medicine (Baltimore).* 2016;95:e4533.
- Wang Z, Zeng J, Li T, Zheng X, Xu X, Ye X, et al. Prevalence of hepatitis B surface antigen (HBsAg) in a blood donor population born prior to and after implementation of universal HBV vaccination in Shenzhen, China. *BMC Infect Dis.* 2016;16:498.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.